



# BioGuano 6000UFP

**BioGuano™ 6000UFP** has all the benefits of BioGuano™ liquid fertiliser in a dry, ultra-fine powder for on-farm mixing and application. Reduced packaging and more concentrated to improve the economics of using Guano fertiliser.

#### Available Sizes:

• 25 kg Kraft Bag	48 per pallet
• 500 kg Bulk Bag	2 per pallet

#### Typical Analysis

##### % w/w

Calcium (as Phosphate).....	30.3%
Phosphorus .....	15.7%
Citrate Soluble .....	6.18%
Water Soluble .....	0.71%
Citrate Insoluble .....	8.88%
Total Organic Carbon .....	1.34%
Magnesium .....	2.34%
Potassium .....	0.19%
Sulphur .....	0.39%
Sodium .....	0.26%
Nitrogen (Organic) .....	0.04%

#### Trace Minerals

##### ppm w/w

Silicon (available) .....	155
Iron .....	1787
Manganese .....	404
Zinc .....	117
Boron .....	15
Copper .....	6
Cobalt .....	1
Molybdenum .....	0.16

Particle Size (average) .....	5 µm
Bulk Density .....	700-750 kg/m³
pH .....	7.45

#### APPLICATION RATES

##### Fertigation:

8-10 kg/ha as required. Application frequency may change depending on the crop and soil fertility.

##### Soil Ameliorant:

5-10 kg/ha as required.

##### Liquid Inject:

2-5 kg/ha. Use a minimum dilution rate of 1 part product to 3 parts water. Application rate will depend on crop, row spacing and soil type. Always consider current soil test results when selecting a crop starter application rate.

##### Foliar Spray:

##### Broadacre Crops, Cereals & Grain Legumes, Lucerne & Pastures:

2-5 kg/ha in 60-120 litres of water depending on canopy development at time of application. Repeat this application as often as required ideally after monitoring soil or leaf nutrient levels.

##### Vegetables:

1 kg per 100 litres water up to a maximum of 7 kg/ha.

##### Orchards, Berries & Vineyards:

1 kg per 100 litres water up to a maximum of 7 kg/ha.

##### Ornamentals & Turf:

1 kg per 100 litres water up to a maximum of 7 kg/ha.

##### Spot-spraying Applications for Small Areas and Newly-Planted Trees, etc:

2-3 g/litre of water to cover 1 m². Can be applied as a foliar and soil-directed spray at label rates with complete safety to crops. Repeat every 2-4 weeks or as required.

##### Storage & Handling:

Store out of direct sunlight in a cool, dry area. Use soon after mixing with water and DO NOT store in the diluted form.

##### Caution:

Always lift with care to avoid personal injury.

DO NOT breathe, swallow or ingest the powder. The dust and diluted spray from this product may act as an irritant. Avoid inhalation and contact with eyes and skin.

Avoid windy conditions when adding product to the tank.

